

A photograph of two men in a gym setting, both wearing athletic wear and jumping rope. The man in the foreground is wearing a light blue long-sleeved shirt and dark shorts, while the man in the background is wearing a grey long-sleeved shirt and dark pants. They are in a bright room with large windows on the left and a wooden shelf with exercise equipment in the background.

ALTUVIIIIO®
Antihemophilic Factor (Recombinant),
Fc-VWF-XTEN Fusion Protein-ehtl

FACTOR UP

INFUSION OVERVIEW

INDICATION

ALTUVIIIIO® [antihemophilic factor (recombinant), Fc-VWF-XTEN fusion protein-ehtl] is an injectable medicine that is used to control and reduce the number of bleeding episodes in people with hemophilia A (congenital Factor VIII deficiency).

Your healthcare provider may give you ALTUVIIIIO when you have surgery.

SELECT IMPORTANT SAFETY INFORMATION

Who should not use ALTUVIIIIO?

You should not use ALTUVIIIIO if you have had an allergic reaction to it in the past.

Please see full [Prescribing Information](#), including Instructions for Use, and Important Safety Information on the last page.

Hypothetical patient

Freedom and flexibility to meet your needs

Ways to treat with ALTUVIIIIO®

ROUTINE PROPHYLAXIS

Prevent bleeds or reduce their frequency with one infusion, once a week

ON-DEMAND TREATMENT

If a breakthrough bleed occurs, it can be treated with as little as one infusion

PERIOPERATIVE MANAGEMENT

Increase your bleed protection during and around the time of surgery

When storing ALTUVIIIIO prior to reconstitution, keep it:

- in the original package to protect it from light
- refrigerated at 2°C to 8°C (36°F to 46°F); do not freeze
- or store it at room temperature, if needed, for up to 6 months. Do not let it exceed 30°C (86°F), and use within the expiration date. Do not return the product to the refrigerator

One way to infuse to fit your needs

Expect the same infusion process, whether it's weekly prophylaxis use, on-demand bleed control, or prior and post-surgery use

SELECT IMPORTANT SAFETY INFORMATION (CONT'D)

What is the most important information I need to know about ALTUVIIIIO? (cont'd)

Do not attempt to give yourself an injection unless you have been taught how by your healthcare provider or hemophilia center. You must carefully follow your healthcare provider's instructions regarding the dose and schedule for injecting ALTUVIIIIO so that your treatment will work best for you.

Please see full [Prescribing Information](#), including Instructions for Use, and Important Safety Information on the last page.



Ready to infuse?

Infuse with confidence! Use the reminder information below any time you need a refresher, once you've been trained on infusions by a healthcare provider. Remember to always review the Instructions for Use before infusing and reach out to your healthcare provider with any questions.



SETTING UP

- Gather medication and supplies
- Double-check the expiration date on the vial to be sure it's correct. Reach out to your healthcare provider if you have any questions about your dosage



INFUSING

- Wrap the tourniquet above where you plan to infuse and clean the site with an alcohol wipe
- Follow the Administration Steps in the Instructions for Use to slowly administer the solution over 1-10 minutes, based on your comfort level.



PREPARING FOR INFUSION

- Wash hands and prepare a clean work surface
- Give medicine time to come to room temperature, then reconstitute the powdered ALTUVIIIIO® with the diluent solution as directed
- Administer reconstituted ALTUVIIIIO as soon as possible. Reconstituted solution may be kept at room temperature, away from sunlight, for up to 3 hours



FINISHING UP

- Once you've infused the full dose, slowly remove the needle
- Throw away all used and unused medical supplies, including empty vials and unused solution, in the appropriate waste container—and you're done!

QUICK TIP



Infuse the same day each week prophylactically, so you never miss a dose—even when traveling

Please see full [Prescribing Information](#), including Instructions for Use, and Important Safety Information on the last page.

LET'S GET STARTED

Watch the [How to infuse ALTUVIIIIO®](#) video for step-by-step instructions.

IMPORTANT SAFETY INFORMATION

What is the most important information I need to know about ALTUVIIIIO?

Do not attempt to give yourself an injection unless you have been taught how by your healthcare provider or hemophilia center. You must carefully follow your healthcare provider's instructions regarding the dose and schedule for injecting ALTUVIIIIO so that your treatment will work best for you.

Who should not use ALTUVIIIIO?

You should not use ALTUVIIIIO if you have had an allergic reaction to it in the past.

What should I tell my healthcare provider before using ALTUVIIIIO?

Tell your healthcare provider if you have had any medical problems, take any medications, including prescription and non-prescription medicines, supplements, or herbal medicines, are breastfeeding, or are pregnant or planning to become pregnant.

What are the possible side effects of ALTUVIIIIO?

You can have an allergic reaction to ALTUVIIIIO. Call your healthcare provider or emergency department right away if you have any of the following symptoms: difficulty breathing, chest tightness, swelling of the face, rash, or hives.

Your body can also make antibodies called "inhibitors" against ALTUVIIIIO. This can stop ALTUVIIIIO from working properly. Your healthcare provider may give you blood tests to check for inhibitors.

The common side effects of ALTUVIIIIO are headache and joint pain.

These are not the only possible side effects of ALTUVIIIIO. Tell your healthcare provider about any side effect that bothers you or does not go away.

Please see full [Prescribing Information](#).

The Sanofi logo, consisting of the word "sanofi" in a lowercase, sans-serif font.

© 2024 Sanofi. All rights reserved.

ALTUVIIIIO and Sanofi are registered trademarks of Sanofi or an affiliate.

MAT-US-2209729-v2.0-06/2024